

Perseus Proteomics Inc. 30-1 Nihonbashi-hakozakicho, Chuo-ku, Tokyo 103-0015, JAPAN

TEL: +81-3-6264-8268 FAX: +81-3-3668-7776 https://www.ppmx.com order@ppmx.com

Anti human SF-1 mouse monoclonal antibody (ProteinA purified)

SF-1: Steroidogenic factor 1

Description

Nomenclature NR5A1

1176200

Code No	PP-N1665-0C
Clone No.	N1665
Lot.	B-2
Concentration	1 mg/mL
Volume	100 uL
lg Class	G1


SF-1 (AD4BP, FTZF1,NR5A1) is still considered as an orphan nuclear receptor belongs to subfamily 5, It was found as a regulator of steroidogenic enzyme gene expression. Oxysterols are suggested as ligands. Its expressed in all sterodogenic tissues, including adrenal cortex, testicular Sertoli cells and Leydig cells, ovarian theca, hypothalamus and anterior pituitary. SF1 plays an important role in adrenal and gonadal development, such as hypothalamic-pituitary-gonadal axis and sex determination.


Application / Recommended Concentration

In order to obtain the best results, optimal working dilutions should be determined by each individual user.

Western Blot	3 ug/mL
Non reducing Western Blot	3 ug/mL
ELISA	0.1 ug/mL (A450=1.0)
Immunoprecipitation	Decide by use
Supershift Assay	Not yet tested
Chromatin immunoprecipitatic	Not yet tested

Immunohistochemistry 10 ug/mL





Rat Adrenal gland paraffin section

Rat Salivary gland paraffin section

Storage

Store at 2 - 8 °C up to one month. For long-term storage, the solution may be frozen in working aliquots. Repeated freezing and thawing is not recommended. Storage in a frost-free freezer is not recommended.

Reference

Suh JM, *et al.* Mol Endocrinol. 2006; 20(12): 3412-20 Qin J, *et al.* Dev Dyn. 2007; 236(3): 810-20

Genbank	076388
Origin	Produced in serum-free medium with hybridoma of mouse myeloma cells (NS-1) and spleen cells derived from a BALB/c mouse immunized with E.coli-expressed recombinant human SF-1(218-461aa)
Specificity	This antibody specifically recognizes human SF-1 and cross reacts with mouse and rat SF-1.
Purification	Affinity chromatography with Protein A
Formulation	Physiologic saline with 0.1 % NaN3.

Notes

Sodium azide may react with lead and copper plumbing to form explosive metal azides. Flush with large amounts of water during disposal.

FOR RESEARCH ONLY. NOT FOR USE IN HUMANS.