

Perseus Proteomics Inc.

In order to obtain the best results, optimal working dilutions should be

4-7-6, Komaba, Meguro-ku, Tokyo 153-0041

Application / Recommended Concentration

determined by each individual user.

TEL: +81-3-5738-1705 FAX: +81-3-3481-5760 https://www.ppmx.com info@ppmx.com

## Anti mouse G9a /Ehmt2 mouse monoclonal antibody

G9a: Euchromatic histone-lysine N-methyltransferase 2

Code No	PP-A8620A-00
Clone No.	A8620A
Lot.	A-2
Concentration	1 mg/mL
Volume	100 uL
lg Class	G2a
Description	Histone H3 Lys 9 (H3-K9) methylation is a crucial epigenetic mark for transcriptional silencing. G9a is the major mammalian H3-K9 methyltransferase that targets euchromatic regions and is essential for murine embryogenesis.

Western Blot	0.5 ug/mL
Non reducing Western Blot	Not yet tested
ELISA	0.1 ug/mL
Immunoprecipitation	Decide by use
Supershift Assay	Not yet tested
Chromatin immunoprecipitatic	Not yet tested
mmunohistochemistry	5 ug/mL
WT mouse ES cell (Positive)	G9a KO mouse ES cell (Negative)

Genbank	AB077209
Origin	Produced in BALB/c mouse ascites after inoculation with hybridoma of mouse myeloma cells (NS-1) and spleen cells derived from a BALB/c mouse immunized with Baculovirus-expressed recombinant mouse G9a (15-114aa).
Epitope	
Specificity	This antibody specifically recognizes mouse G9a and cross reacts with human G9a. Not yet tested in other species.

Store at 2 - 8 °C up to one month. For long-term storage, the solution may be frozen in working aliquots. Repeated freezing and thawing is not recommended. Storage in a frost-free freezer is not recommended.

Purification Ammonium sulfate fractionation. Physiologic saline with 0.1 % NaN3. Formulation

Tachibana M. et al. Genes Dev, 2005; 19(7): 815-26 Reference

Notes

Sodium azide may react with lead and copper plumbing to form explosive metal azides. Flush with large amounts of water during disposal.

## FOR RESEARCH ONLY. NOT FOR USE IN HUMANS.